

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 900 841 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

10.03.1999 Bulletin 1999/10

(51) Int. Cl.⁶: **C12N 15/12**, C07K 14/725,

C07K 16/28, A61K 38/17,

A61K 39/395, G01N 33/53

(21) Application number: 97401404.5

(22) Date of filing: 18.06.1997

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

Designated Extension States:

AL LT LV RO SI

(71) Applicants:

- INSTITUT NATIONAL DE LA SANTE ET DE LA
RECHERCHE MEDICALE (INSERM)
75654 Paris Cédex 13 (FR)
- INSTITUT GUSTAVE ROUSSY
F-94805 Villejuif Cédex (FR)
- APPLIED RESEARCH SYSTEMS ARS HOLDING
N.V.
Curacao (AN)

(72) Inventors:

- Trlebel, Frédéric
78000 Versailles (FR)
- Mastrangeli, Renato
00146 Rome (IT)
- Romagnani, Sergio
50100 Firenze (IT)

(74) Representative:

Hirsch, Denise et al
Cabinet Lavoix
2, Place d'Estienne d'Orves
75441 Paris Cedex 09 (FR)

(54) **LAG-3 splice variants**

(57) This invention concerns an isolated nucleotidic
sequence selected from the group consisting of :

- a) the nucleotidic sequences SEQ ID N° 1, SEQ ID
N°3 or SEQ ID N°5 ;
- b) the nucleotidic sequences which hybridize under
stringent conditions to any of the sequences
defined in a) and which code for a polypeptide
which is a variant of the LAG-3 molecule ;
- c) the nucleotidic sequences which are degener-
ated as results of the genetic code to the nucleotidic
sequences defined in a) and b), and which code for
a polypeptide which is a genetic variant of the LAG-
3 molecule.

EP 0 900 841 A1

Description

[0001] The present invention relates to the identification of differentially spliced variants of the LAG-3 molecule and their use as immunomodulators.

5 [0002] It is now recognized that the proteins encoded by MHC Class II region are involved in many aspects of immune recognition, including the interaction between different lymphoid cells such as lymphocytes and antigen presenting cells. Different observations have also shown that other mechanisms which do not take place via CD4 participate in the effector function of T helper lymphocytes.

[0003] The lymphocyte activation gene 3 (LAG-3) expressed in human activated T and NK cells encodes a 498 amino-
 10 acids (aa) type I membrane protein with four extracellular immunoglobulin superfamily (IgSF) domains (1). Analysis of this sequence revealed notable patches of identity with stretches of aa sequences found at the corresponding positions in CD4, although the overall aa sequence homology with human CD4 is barely above background level (approximately 20 % sequence identity). There are also some internal sequence homologies in the LAG-3 molecule between domains 1 (D1) and 3 (D3) as well as between domains 2 (D2) and 4 (D4) suggesting that LAG-3 has evolved like CD4 by gene
 15 duplication from a preexisting 2 IgSF structure (1). In addition, LAG-3 and CD4 genes are located in a very close proximity on the distal part of the short arm of chromosome 12 (2, B. Huard, P. Gaulard, F. Faure, T. Hercend, F. Triebel, Immunogenetics 39, 213, 1994). LAG-3 and CD4 can therefore be regarded as evolutionary "first cousins" within the IgSF (2).

[0004] Using a quantitative cellular adhesion assay, the authors of the present invention previously showed that
 20 rosette formation between LAG-3 transfected Cos-7 cells and MHC class II⁺ B lymphocytes was specifically dependent on LAG-3/MHC class II interaction (3). A direct and specific binding of LAG-3 to various human class II molecules (including different alleles and isotypes), as well as to murine and monkey class II molecules has also been observed with a LAG-3Ig fusion protein (4). This dimeric LAG-3Ig recombinant globulin binds MHC class II monomorphic residues with a much higher avidity ($K_d = 60$ nM at 37° C) than CD4Ig (5); LAG-3Ig is indeed able to block CD4/MHC class II interaction in an intercellular adhesion assay (5).
 25

[0005] The role of LAG-3/class II molecule interaction has been investigated using LAG-3 specific monoclonal antibodies (mAb) (6) and LAG-3Ig molecules (7). This interaction leads to downregulation of T cell clone activation. Productive LAG-3/MHC class II interaction is mediated through T-T cell contacts, presumably via negative MHC class II signaling into T cells. Overall, LAG-3 is expressed only after lymphocyte activation *in vitro* as well as *in vivo* (6) and
 30 hence does not play a role in the induction phase of the response, in contrast to CD4. In addition, mAb blocking experiments have shown that LAG-3 does not participate in the recognition phase of MHC class II-restricted CD4⁺ T cell clones. The functional role of LAG-3 is therefore strikingly different from that of the other MHC ligands, CD4 and CD8. It is currently thought that T cell MHC class II molecules have a role similar to CTLA-4 following LAG-3 binding (4), i.e. induction of clonal deletion of previously activated T cells (8).

35 [0006] More recently, LAG-3 has been found to be preferentially expressed and released by activated Th1 cells, i.e. T helper cells producing IFN- γ and TNF, and to be upregulated by IL-12, a powerful Th1-inducing cytokine (F. Annunziato, R. Manetti, L. Tomasovic, MG. Giudizi, R. Biagiotti, V. Giannò, P. Germano, C. Mavilia, E. Maggi and S. Romagnani, FASEB J. 10, 769-775, 1996). High levels of soluble LAG-3 (sLAG-3) were also found in the serum of patients with relapsing multiple sclerosis (MS) (ibidem) and of a few patients with systemic lupus erythematosus (S. Romagnani, Clin. Immunol. Immunopath. 80, 225-235, 1996). These findings suggest that LAG-3 could represent an useful diagnostic
 40 marker for Th1-mediated immune diseases, such as Hashimoto's thyroiditis, type I diabetes mellitus, multiple sclerosis, Chron's disease, rheumatoid arthritis, acute allograft rejection and acute graft-versus-host disease (GVHD). On the other hand, the presence of high levels of sLAG-3 in the serum of MS patients and possibly of other patients suffering of the above diseases, supports a naturally occurring protective role for sLAG-3, as it could compete with the membrane-bound form and thus block the LAG-3-mediated immune responses.
 45

[0007] The authors of the present invention have investigated whether other forms of LAG-3 are expressed, due to alternative splicing of the nuclear transcript.

[0008] Most eukaryotic protein encoding genes contain the sequences present in the corresponding mature mRNA in discontinuous DNA fragments (exons) interspersed among sequences (introns) that do not form a part of the mature
 50 mRNA. The primary transcripts of these genes thus contain both sequences corresponding to exons and sequences corresponding to introns. The intron sequences are subsequently excised by a nuclear multistep process, known as pre-mRNA splicing. Namely, introns are demarcated by consensus sequences at their 5'(donor) and 3'(acceptor) boundaries. The splicing process comprises cleavage at the acceptor site with concomitant ligation of the 5' and 3' exons.

55 [0009] In most cases, the exons present in a gene are incorporated into one mature mRNA through an invariant ligation of consecutive pairs of donor and acceptor splice sites, providing a single gene product.

[0010] In some cases however, the same gene contain alternate splice sites which may lead to different transcripts. The mechanism regulating this alternative splicing are poorly understood, but at least for some genes, it is known to be

cell and developmental specific.

[0011] The alternative splicing leads to the production of multiple protein isoforms from a single gene. It is one of the molecular process responsible for generating the protein diversity.

[0012] More than 50 genes are currently known to generate protein diversity through the use of alternative splicing.

5 This mechanism is particularly common and elaborated among the contractile protein genes.

[0013] The authors of the present invention have discovered three new differentially spliced LAG-3 variants, respectively referred as LAG-3V1, LAG-3V2, and LAG-3V3.

[0014] LAG-3V1 encodes a soluble 36 kD protein containing 8 new aminoacid residues after the D2 domain, LAG-3V2 encodes a transmembrane protein of 61 kD which does not contain domain D4, and LAG-3V3 encodes a soluble 52 kD protein containing 8 new aminoacid residues after domain D3.

10 [0015] These variants provide new approaches for studying both the regulation of the LAG-3 gene and the biological function of the protein, and are useful for the manufacture of new immunomodulator compounds, especially compounds which mimic the biological function of LAG-3 or which can act as agonists or antagonists of the interaction between LAG-3 and the MHC class II molecules.

15 [0016] The subject of the present invention is thus an isolated nucleotidic sequence selected from the group consisting of:

a) the nucleotidic sequences SEQ ID N° 1, SEQ ID N° 3 or SEQ ID N° 5;

20 b) the nucleotidic sequences which hybridize under stringent conditions to any of the sequences defined in a) and which code for a polypeptide which is a variant of the LAG-3 molecule;

c) the nucleotidic sequences which are degenerated as results of the genetic code to the nucleotidic sequences defined in a) and b), and which code for a polypeptide which is a genetic variant of the LAG-3 molecule.

[0017] The present invention is also directed to the purified polypeptides encoded by the nucleotidic sequences defined above. Said polypeptides are designated hereinafter as "LAG-3 variants". The present invention is also relating to pharmaceutical compositions comprising a LAG-3 variant polypeptide. These compositions are useful for treating immune-related pathologies, in particular Th1-dependent diseases such as Hashimoto's thyroiditis, type I diabetes mellitus, multiple sclerosis, Chron's disease, rheumatoid arthritis, acute allograft rejection, acute GVHD, Grave's ophtalmopathy, cerebral malaria, Lyme arthritis, reactive arthritis (Yersinia-induced), HCV-induced chronic hepatitis, primary sclerosing colangitis, contact dermatitis, unexplained recurrent abortion, aplastic anaemia, and Helicobacter pilori-induced gastric antritis.

[0018] The present invention also relates to the use of the LAG-3 variants for the manufacture of immunomodulator compounds. Such compounds can mimic or alter the biological function of LAG-3 and/or LAG-3 variants, inducing therefore some modifications of the cellular interactions involving the participation of LAG-3 or its variants.

35 [0019] The present invention is also directed to poly- or monoclonal antibodies, Fab, Fab' and F(ab')₂ or Fv fragments thereof, directed to a specific epitope of one of the above-defined LAG-3 variants. The use of these specific antibodies for purifying said variants or for preparing therapeutic or diagnostic compositions, as well as the resulting compositions, are also comprised within the scope of the present invention.

[0020] The present invention further provides a therapeutic method for treating immune-related pathologies comprising the administration to a patient of a LAG-3 variant in an efficient quantity or a composition comprising said variant as an active ingredient.

[0021] The invention is also relating to a therapeutic method for treating immune-related pathologies comprising the administration to a patient of an anti-LAG-3 variant- antibody as defined here-above, or a therapeutic composition containing said antibody as active ingredient.

45 [0022] In a preferred embodiment, the invention is directed to an isolated nucleotide sequence selected from the group consisting of SEQ ID N° 1, SEQ ID N° 3, SEQ ID N° 5 or the fully complementary sequences thereof.

[0023] The invention also relates to a purified polypeptide which is resulting from the expression of one of the nucleotidic sequences of the invention. Preferably, the polypeptide is selected from the group consisting of LAG-3V1, LAG-3V2 and LAG-3V3, respectively encoded by the sequences SEQ ID N° 1, SEQ ID N° 3 and SEQ ID N° 5 and respectively corresponding to the sequences SEQ ID N° 2, SEQ ID N° 4 and SEQ ID N° 6.

50 [0024] The polypeptides according to the invention can be obtained by any of the standard methods of purification of membrane or soluble proteins, by peptide synthesis or by application of genetic engineering techniques. Said techniques comprise the insertion of a nucleotidic sequence coding for one of the peptide of the invention into an expression vector, such as a plasmid, and the transformation of host cells with this expression vector, by any of the methods available to the skilled person, like for instance electroporation.

55 [0025] Advantageously, the nucleotidic sequences of the invention are obtained by RT-PCR (Reverse Transcriptase Polymerase Chain Reaction) of the corresponding LAG-3 variant cDNA. Briefly, total RNA is extracted from activated human peripheral blood lymphocytes (PBL), and is reverse-transcribed and amplified using a specific pair of oligonu-

cleotides primers. In the case of the present invention, the primers are consisting of a sense primer and an antisense primer which are located at two specific positions on the LAG-3 sequence which will be described in more details in the following examples. The resulting cDNA is then amplified by enzymatic amplification and can be subcloned in an appropriate host cell.

- 5 [0026] The present invention also relates to expression vectors comprising a nucleotidic sequence coding for a polypeptide according to the invention and host cells transformed with these vectors.
- [0027] An "expression vector" refers to a replicable DNA construct used either to amplify or to express DNA which encodes one of the polypeptides of the invention.
- 10 [0028] Host cells may be prokaryotic or eukaryotic, including but not limited to bacteria, yeasts, insect cells, mammalian cells, including cell lines which are commercially available.
- [0029] The present invention is also directed to anti-LAG-3 variants antibodies or fragments thereof, as defined above. Optionally, these antibodies or their fragments can be linked to a marker (radioisotope, fluorescent dye, enzymatic marker...) or to a therapeutically active molecule, which is for example a cytotoxic compound.
- 15 [0030] The polyclonal antibodies may be prepared according to well-known methods, such as that described by BEN-EDICT A.A. et al. (). Methods of production of monoclonal antibodies are well known from the prior art, especially the one described by KOHLER and MILSTEIN. This method, together with variants thereof, are described by YELTON et al. ().
- [0031] These antibodies can be used in a method for purifying a polypeptide according to the invention, or for a dosage or identification method. Said method is for example, but not limited to, a radio-immunological method of the RIA or
- 20 IRMA type, an immuno-enzymatic method, like the ELISA method, or a direct or indirect immunofluorescence method.
- [0032] The invention also includes the use of said antibodies for the manufacture of therapeutic compositions able to block the activity of LAG-3 variants, i.e. having immunomodulatory activities, such as induction of the maturation, differentiation, proliferation and/or function of cells expressing a LAG-3 variant, e.g. activated T and NK cells.
- [0033] The antibodies to LAG-3 variants may be used as potentiators of vaccines or immunostimulants in immunosuppressed patients, such as patients infected with HIV or treated with immunosuppressant substances.
- 25 [0034] The therapeutic compositions according to the present invention comprise soluble LAG-3 variant proteins or antibodies as defined above, as well as a pharmaceutically acceptable vehicle. These compositions may be formulated according to the usual techniques. The vehicle can vary in form in accordance with the chosen administration route: oral, parenteral, sublingual, rectal or nasal.
- 30 [0035] For the compositions for parenteral administration, the vehicle will generally comprise sterile water as well as other possible ingredients promoting the solubility of the composition or its ability to be stored. The parenteral administration route can consist of intravenous, intramuscular or subcutaneous injections.
- [0036] The therapeutic composition can be of the sustained-release type, in particular for long-term treatments, for example in autoimmune diseases. The dose to be administered depends on the subject to be treated, in particular on
- 35 the capacity of his/her immune system to achieve the desired degree of protection. The precise amounts of active ingredient to be administered may be readily determined by the practitioner who will initiate the treatment.
- [0037] The therapeutic compositions according to the invention can comprise, in addition to soluble LAG-3 variants or the antibodies according to the invention, another active ingredient, where appropriate, bound via a chemical bond to LAG-3 variant or to an antibody according to the invention. As an example, there may be mentioned soluble LAG-3
- 40 variant proteins according to the invention fused to a toxin, for example ricin or diphtheria anatoxin, capable of binding to MHC Class II molecules and of killing the target cells, for example leukaemic or melanoma cells, or fused to a radioisotope.
- [0038] The examples which follow, together with the attached reference figures, will illustrate the invention in greater detail.

45

LEGENDS TO FIGURES

[0039]

- 50 - Fig. 1 represents the schematic structure of wild type (wt) LAG-3 and its variants V1, V2 and V3. LAG-3V1 derives from the retention of intron 4, i.e. cleavage at the donor and acceptor sites flanking intron 4 does not occur. An in-frame stop codon located after 8 codons in the retained intron 4 leads to a truncated soluble LAG-3V1 protein, containing D1, D2 and 8 new amino acid residues.
- 55 LAG-3V2 lacks exon 6, due to the ligation of the donor site on intron 5 to the acceptor site on intron 6. As no shift of the reading frame occurs, the resulting LAG-3V2 protein is a transmembrane protein which does not contain the D4 domain.
- LAG-3V3 derives from cleavage of the nuclear transcript at a different polyadenylation site located ~170 bp

downstream to the 5' end of intron 5. The retained intron 5 sequence contains an in-frame stop codon. The resulting mRNA encodes a truncated soluble protein which contains D1, D2, D3 and 8 new amino acid residues.

- 5 - Fig. 2 is a schematic representation of the intron/exon organization of the LAG-3 gene. Splicing events which result in the generation of RNA transcripts encoding either wtLAG-3, LAG-3V1, LAG-3V2 or LAG-3V3 are indicated by dotted lines.

SP: signal peptide; D1-D4: IgSF domains 1-4; TM: transmembrane sequence; CYT: cytoplasmic domain; I1-I9: intron 1-9.

10 New exons in LAG-3V1 and LAG-3V3 are shown as shaded boxes.

- Fig. 3 is a schematic representation of wtLAG-3 and LAG-3V1. The location of the primers and probes used for the isolation of LAG-3V1 is indicated by arrows. DNA fragments amplified using the specific primers are shown by single lines. The size of the fragments obtained by RT-PCR is indicated.
- 15 - Fig. 4 shows a Southern blot analysis of wtLAG-3 and LAG-3V1 cDNA fragments amplified by RT-PCR with primers F459 and R460.

- 20 a) ethidium bromide stained gel. b) blot hybridized with a LAG-3 specific cDNA probe. The lower band derives from wtLAG-3 while the upper one, evident only after hybridization, derives from LAG-3V1.

- Fig. 5 shows a Southern blot analysis of LAG-3V1 cDNA fragments amplified by RT-PCR with primers F176 and R460. a) ethidium bromide stained gel. The major band derives from wtLAG-3. b) blot hybridized with the LAG-3 intron 4 specific oligoprobe I4. The upper band derives from LAG-3V1 while the lower one is a wtLAG-3/LAG-3V1 heteroduplex.
- 25

- Fig. 6 is a schematic representation of wtLAG-3, LAG-3V2 and LAG-3V3. The location of the primers and probes used for the isolation of LAG-3V2 and LAG-3V3 is indicated by arrows. DNA fragments amplified using the specific primers are shown by single lines. The size of the fragments obtained by RT-PCR is indicated.

- 30 - Fig. 7 shows a Southern blot analysis of wtLAG-3, LAG-3V2 and LAG-3V3 cDNA fragments amplified by RT-PCR with primers F176 and R401.

- a) ethidium bromide stained gel. b) blot hybridized with the LAG-3D2 specific oligoprobe F459. The upper band derives from wtLAG-3 while the 780 and 940 bp bands derive from LAG-3V2 and LAG-3V3, respectively.

- 35 - Fig. 8 shows the agarose gel electrophoresis pattern of RT-PCR products obtained with LAG-3V2 and LAG-3V3 specific primers, i.e. 173/V2R and 173/V3R, respectively.

- Fig. 9 shows the Western blotting patterns of PHA-blasts with anti LAG-3 in mAbs.

EXAMPLES

Example I:

Cloning of human LAG-3 variant 1 (LAG-3V1) by RT-PCR on human PBMC (Peripheral Blood Mononuclear Cells)

45 - RNA extraction and RT-PCR:

[0040] PBMC were isolated by Ficoll-Hypaque density gradient centrifugation and activated with 1 µg/ml PHA and 100 U/ml IL-2 for 48 hr at 37°C, 5% CO₂. mRNA was extracted by the Oligotex Direct mRNA kit (Quiagen Inc., Chatsworth, CA, USA.) and reverse-transcribed with an oligo(dT) primer, using a first strand synthesis kit (RT-PCR kit, Stratagen, La Jolla, CA, USA).

50 [0041] Amplification of cDNA was performed with 2.5 U Taq polymerase and 100 ng each of forward primer F459 (5' TCTCTCAGAGCCTCCGATGGGTCAATTTG 3') (SEQ ID N° 7) and reverse primer R460 (5' TCCTGCAGATGGATATGGCAGGTGTAGGTC 3') (SEQ ID N° 8) which anneal to nucleotides 762-791 and 1217-1246 of LAG-3 sequence, respectively (Fig. 3). Thirty PCR cycles were performed, each cycle including denaturation at 94°C for 1 min, annealing at 66°C for 1 min, and extension at 72°C for 2 min.

- Analysis of amplified DNA by Southern blotting :

[0042] A 10 µl aliquot of the amplified DNA was fractionated on a 2 % agarose gel. The expected 485 bp LAG-3 fragment was observed (Fig. 4a). After blotting onto nitrocellulose membrane filter and hybridization with a ³²P-labelled LAG-3D1D2 specific cDNA probe obtained by PCR, in addition to the 485 bp band a ~890 bp band was found. The 890 bp fragment was reamplified, cloned and sequenced.

- Cloning and sequencing of the 890 bp LAG-3 fragment:

[0043] The 890 bp fragment was cloned into the vector pCR™II (Promega) and the insert was sequenced using the ABI Prism Dye Terminator Cycle Sequencing Ready Reaction Kit (Perkin-Elmer) and the automated DNA sequencer mod. 373A (Perkin-Elmer). The results indicate that this fragment is 892 bp in length and derives from a LAG-3 variant retaining intron 4 (Fig. 2). Due to the presence of an in-frame stop codon in intron 4, this variant, named LAG-3V1, is predicted to encode a soluble 36 kD protein containing domains D1, D2 and 8 new C-terminal amino acid residues (Fig. 1).

- Confirmation of LAG-3V1 expression :

[0044] To confirm that the LAG-3V1 fragment derives from mRNA and not from genomic DNA, a second RT-PCR experiment was performed using the forward primer F176 (5' CCTGGGCCAGGCCTCGA TGAC 3') (SEQ ID N° 9) which anneals to nucleotides 725-745 of LAG-3 sequence, and the reverse primer R460 (Fig. 3). Primer F176 which spans the D1/D2 splice junction site, should not allow the amplification of LAG-3 genomic DNA.

[0045] Upon agarose gel electrophoresis of the PCR products, the expected ~520 bp wtLAG-3 band was observed (Fig. 5a). Southern blotting with the LAG-3 intron 4 specific oligoprobe I4 (5' CCCCACTCTGCTTCACATT 3') (SEQ ID N° 10) showed the expected 930 bp LAG-3V1 band and an additional ~800 bp band (Fig. 5b). The two fragments were reamplified and sequenced. The upper band was confirmed to correspond to LAG-3V1 while the lower one was a wtLAG-3/LAG-3V1 heteroduplex.

Example II:

Cloning of human LAG-3 variant 2 (LAG-3V2) and LAG-3 variant 3 (LAG-3V3) by RT-PCR on activated PBMC

- RNA extraction and RT-PCR :

[0046] PBMC were isolated by Ficoll-Hypaque density gradient centrifugation and activated with 1 µg/ml PHA and 100 U/ml IL-2 for 48 hr at 37°C, 5% CO₂.

[0047] Total RNA was extracted by TriZol reagent (GIBCO-BRL). Total RNA (5µg) was reverse-transcribed using the anchor oligo(dT) primer mixture AT17A, AT17C and AT17G (5' GGCCCTGGATCCGGACCTAA(T)₁₇ (SEQ ID N° 1) followed by A, C or G) (Fig. 6) which allows to obtain cDNA molecules with homogeneous 3' ends and an anchor for subsequent PCR.

[0048] Reverse transcriptase (RT) reaction was performed in 50 µl final volume of 1x RT buffer, 10 mM DTT, 1 mM dNTP, 40 U RNase inhibitor (Boehringer) and 400 U Superscript II RT (GIBCO-BRL) at 37°C for 90 min. RT reaction was stopped at 90°C for 5 min. RNA was digested with 1 U RNase H (Boehringer) at 37°C for 30 min.

[0049] PCR was performed with the forward primer F176 and the reverse primer R401 (5' GGCCCTGGATCCGGACCTAA 3') (SEQ ID N° 12) which anneals to the 3' anchor of cDNA (Fig. 6). Primers F176 and R401 should allow to amplify all LAG-3 splice variants containing the D1D2 junction up to poly(A) tail.

[0050] PCR was performed on 5 µl cDNA equivalent to 0.5 µg total RNA in 100 µl final volume containing 1x Taq buffer, 2.5 U Taq DNA polymerase (Advanced Biotechnologies), 1.5 mM MgCl₂, 200 µM dNTP, 10% DMSO and 50 pmoles each of F176 and R401 primers. A hot start technique and 30 PCR cycles (96°C for 30 sec, 65°C for 30 sec, 72°C for 4 min) were used. RNA without RT was used as negative control in PCR experiment.

- Southern blot analysis of RT-PCR products :

[0051] RT-PCR products (10 µl) were fractionated by agarose gel electrophoresis and blotted onto Hybond N+ nylon membrane (Amersham). The blot was hybridized with the DIG-labeled LAG-3D2 specific oligoprobe F459 (Fig. 6) in 5X SSC, 0.02% SDS, 0.5% blocking reagent (Boehringer), 0.1% N-laurylsarcosine at 55 °C for 30 min.

[0052] The blot was washed twice with 2x SSC, 0.1% SDS at 55°C. Hybrid detection was performed with HRP-conjugated anti-DIG antibodies (Boehringer) and ECL reagents (Amersham).

[0053] The results showed that in addition to the expected 1.12 kb wtLAG-3 fragment two minor bands of approximately 940 and 780 bp were present both in the ethidium bromide stained gel (Fig. 7a) and in the Southern blot (Fig. 7b).

5 - Sequencing of LAG-3 fragments :

[0054] The 940 and 780 bp PCR fragments were isolated from agarose gel, reamplified with F176/R401 primers and directly sequenced. The results showed that the 780 bp fragment derives from an in-frame skipping of LAG-3 exon 6 (Fig. 2). This variant, named LAG-3V2, is predicted to encode a 61 kD transmembrane protein which does not contain the domain D4 (Fig. 1).

[0055] Sequencing of the 940 bp fragment showed that it derives from the cleavage of the nuclear transcript at a different polyadenylation site located ~170 bp downstream to the 5' end of intron 5 (Fig. 2). The retained intron 5 sequence contains an in-frame stop codon. The resulting mRNA encodes a 52 kD soluble LAG-3 variant, named LAG-3V3, which contains domain D1, D2 and D3 followed by 8 new amino acid residues (Fig. 1). Both LAG-3V2 and LAG-3V3 maintain the 4 glycosylation sites of wtLAG-3.

- Confirmation of LAG-3V2 and LAG-3V3 expression :

[0056] To confirm the expression of LAG-3V2 and LAG-3V3 and to assess the presence of the entire 5' sequence encoding the LAG-3 N-terminus, RT-PCR was performed on total RNA from activated PBMC with the forward primer 173 (5' TATAGGATCCGGTGCCCAG ACCATAGGAGAGATG 3') (SEQ ID N° 13) which anneals to nucleotides 212-233 of LAG-3 sequence spanning the ATG start codon, and the reverse primers V2R (5'GGCGTTCACGTGGTTGGGCAC-CTGTGATGATT 3') (SEQ ID N° 14) or V3R (5'TCACCTACTCGAGAAAAGTGGGGGCCGAGAT 3') (SEQ ID N° 15) (Fig. 6). As the primer V2R anneals to the splice junction site D3/TM of LAG-3V2 and the primer V3R anneals to the 5' sequence of intron 5 retained in LAG-3V3, only amplification of these two variants should occur.

[0057] PCR was performed with annealing temperature at 68°C for the first two cycles and 72°C for the remaining 28 cycles. Upon agarose gel electrophoresis, the expected 1.10 Kb LAG-3V2 and 1.23 Kb LAG-3V3 fragments were found (Fig. 8). DNA sequencing of the amplified fragments confirmed that LAG-3V2 encodes a 61 kD transmembrane protein containing domains D1, D2, D3, the transmembrane domain (TM) and the cytoplasmic domain (CYT), while LAG-3V3 encodes a 52 kD soluble protein containing D1, D2 and a new 8 aa C-tail (Fig. 1).

[0058] In conclusion, at present four LAG-3 molecules have been discovered, two membrane-bound (wtLAG-3 and LAG-3V2) and two soluble forms which lack the transmembrane sequence (LAG-3V1 and LAG-3V3). The analysis of LAG-3 variant expression in specific T cell subsets and in different activation states can provide useful information for a better understanding of the LAG-3 function

Example III:

Characterization of LAG-3V3 protein

[0059] Human peripheral blood mononuclear cells (PBMC) were obtained by centrifugation on ficoll-diatrizoate density gradient from buffy coats of blood donation units. PBMC were obtained from two healthy donors. PBMC collected from the interface between the supernatant and the gradient were washed three times in PBS and finally resuspended at the concentration of 2.10^6 cells/ml in RPMI 1640 culture medium enriched with 10 % fetal bovine serum (FBS), 2 mM L-glutamine, penicillin 100 IU/ml and streptomycin 100 µg/ml. Phytohemagglutinin (PHA) and recombinant human interleukin (IL-2) were then added at a final concentration of 1 µg/ml and 100 U/ml respectively.

[0060] The cells were incubated at 37°C in 5 % CO₂ atmosphere for 72 and 120 hours. At the end of the incubation times, cell suspensions were transferred to 50 ml polypropylene tubes and centrifuged at 400 g for 10 minutes. The supernatants were collected and immediately frozen at -20°C. The total protein determination was carried out by the Bradford method, using the Coomassie Plus Protein Assay (Pierce, Rockford, IL, USA). For analysis, 100 µg of samples were completely dried under vacuum centrifugation, resuspended in 10 µl bi-distilled water (ddH₂O), 10 µl sample buffer 2x were added and the samples were incubated 5 minutes at 100°C.

[0061] The samples were then loaded on 8 % polyacrylamide Tris glycine gels for SDS-PAGE analysis and run under denaturing and reducing conditions at a constant 125 volts. The run was stopped when the dye front reached the bottom of the gel.

[0062] The gel was removed and incubated for 20 minutes in transfer buffer. A nitrocellulose membrane 0.45 µm was incubated for 10 minutes in transfer buffer in a separate container with gentle agitation. The proteins in the gel were then transferred electrophoretically to the membrane for 90 minutes at a constant 75 volts. When the transfer was completed, the membrane was dried at room temperature and then incubated for 20 minutes in 1 % (w/v) KOH with gentle agitation

and washed 4 times with PBS for 5 minutes. The membrane was stained 5 minutes in Ponceau S solution to show proteins and molecular weight markers and destained 4 times with PBS. Non-specific binding sites were blocked by incubating the membrane overnight at 4°C with 1 % Mile in PBS/0.5 % Tween. After elimination of blocking solution, the membrane was blotted with anti-LAG-3 monoclonal antibody (mAb) 2H6 diluted at 5 µg/ml blocking solution. After one hour of incubation at room temperature with agitation, the membrane was washed 5 times with PBS/0.5 % Tween for 10 minutes each and than incubated for one hour at room temperature in agitation with goat anti-mouse polyclonal antibodies conjugated with horseradish peroxidase (HRP) diluted 1:1000 in blocking solution. At the end of the incubation, the membrane was washed as described above.

10 RESULTS

[0063] The immunoblotted proteins were then revealed using the enhanced-chemiluminescent (ECL) method (Amersham Italia, Milan Italy).

15 [0064] Lanes 1 and 3 of figure 9 represent the results obtained from donor 1, respectively after 72 and 120 hours incubation ; whereas lanes 2 and 4 correspond to donor 2 and lane 5 to the lymphoblastoid cell line RAJI as negative control. The starting protein concentrations were respectively 3.1 mg/ml; 1.9 mg/ml; 2.8 mg/ml; 3.4 mg/ml; 3.4 mg/ml from lane 1 to lane 5.

20 [0065] A band with an apparent molecular weight of about 5.5 kDa reacted with anti-LAG-3 mAb. This apparent molecular weight is in agreement with that expected for the putative LAG-3 variant 3 based on the mRNA sequence. This band specifically appears in PHA-blasts and is not present in the supernatant of the B-lymphoblastic cell line RAJI, processed as the supernatants from PHA-blasts.

25

30

35

40

45

50

55

SEQUENCE LISTING

5 (1) GENERAL INFORMATION:

(i) APPLICANTS:

(A) NAME: Institut National de la Santé et de la
Recherche Médicale
(B) STREET: 101 rue de Tolbiac
(C) CITY: Paris
(E) COUNTRY: France
10 (F) POSTAL CODE (ZIP): 75013

(A) NAME: Institut Gustave Roussy
(B) STREET: 39 rue Camille Desmoulins
(C) CITY: Villejuif
(E) COUNTRY: France
15 (F) POSTAL CODE (ZIP): 94805
(G) TELEPHONE: 0142114211
(H) TELEFAX: 0142115300

(A) NAME : Applied Research Systems ARS Holding N.V
20 (B) STREET : 6 John R. Gorsiraweg PO Box 3889
(C) CITY : CURACAO
(E) COUNTRY : The Dutch West Indies

25 (ii) TITLE OF INVENTION: Lag-3 variants

(iii) NUMBER OF SEQUENCES: 15

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
30 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2279 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

45 (*i) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TCAGGCTGCC TGATCTGCCC AGCTTTCCAG CTTTCCTCTG GATTCCGGCC TCTGGTCATC 60
CCTCCCCACC CTCTCTCAA GGCCCTCTCC TGGTCTCCCT TCTTCTAGAA CCCCTTCCTC 120
CACCTCCCTC TCTGCAGAAC TTCTCCTTTA CCCCCACCC CCCACCACTG CCCCTTTCC 180
50 TTTTCTGACC TCCTTTTGGA GGGCTCAGCG CTGCCAGAC CATAGGAGAG ATGTGGGAGG 240
CTCAGTTCCT GGGCTTGCTG TTTCTGCAGC CGCTTTGGGT GGCTCCAGTG AAGCCTCTCC 300

55

EP 0 900 841 A1

	AGCCAGGGGC TGAGGTCCCG GTGGTGTGGG CCCAGGAGGG GGCTCCTGCC CAGCTCCCCT	360
	GCAGCCCCAC AATCCCCCTC CAGGATCTCA GCCTTCTGCG AAGAGCAGGG GTCACTTGGC	420
5	AGCATCAGCC AGACAGTGGC CCGCCCGCTG CCGCCCCCGG CCATCCCCTG GCCCCCGGCC	480
	CTCACC CGGC GCGCCCTCC TCCTGGGGGC CCAGGCCCCG CCGCTACACG GTGCTGAGCG	540
	TGGGTCCCGG AGGCCTGCGC AGCGGGAGGC TGCCCTGCA GCGCCGCGTC CAGCTGGATG	600
10	AGCGCGGCGG GCAGCGCGGG GACTTCTCGC TATGGCTGCG CCCAGCCCGG CGCGCGGACG	660
	CCGCGCAGTA CCGCGCCGCG GTGCACCTCA GGGACCGCGC CCTCTCCTGC CGCCTCCGTC	720
	TGCGCCTGGG CCAGGCCTCG ATGACTGCCA GCGCCCCAGG ATCTCTCAGA GCCTCCGACT	780
15	GGGTCATTTT GAAGTGTCC TTCAGCGGCC CTGACCGCCC AGCCTCTGTG CATTGGTTCC	840
	GGAACCGGGG CCAGGGCCGA GTCCCTGTCC GGGAGTCCCC CCATCACCAC TTAGCGGAAA	900
	GCTTCCTCTT CCTGCCCAA GTCAGCCCCA TGGACTCTGG GCCCTGGGGC TGCATCCTCA	960
20	CCTACAGAGA TGGCTTCAAC GTCTCCATCA TGTATAACCT CACTGTTCTG GGTAACCTCC	1020
	CCACTCTGCT TCACATTGA CCACAACTCC TTCCTGCCCC CTTGTACAC TCCCCTAACT	1080
	ATGGGTCCCC AAACCAGGT CTGCGCAGCG AGTGGCCTAC GTCATTGCTG TGGGTCTCAC	1140
	TGTTGACCC CTTTATATTG CTGGCAGCCT CACAGCTGCC ATCACCCTT CTGCTTCTC	1200
25	CCGTGGCCTT CCAGCGTCAT TGCCGCGCTT CCCTCTCCTT CCGACTAAGC CCACTTGCTG	1260
	GGTTTCTGAG CCTCCTCAGC TCATCACCTT ATTCTGCTCC TTAGCACTCT TATGAGCCAG	1320
	ACCATCTCCT GAATTCTTCT GCCTCCCTTC CTTGCAGCCC CAGCACTCCC TCCCCACTGC	1380
30	AGCACCAGC TTTAACTTTG GGTTTTCTTT TCTCTTCAGG TCTGGAGCCC CCAACTCCCT	1440
	TGACAGTGA CGCTGGAGCA GGTTCAGGG TGGGGCTGCC CTGCCGCTG CCTGCTGGTG	1500
	TGGGGACCG GTCTTCTCTC ACTGCCAAGT GGAATCCTCC TGGGGAGGC CCTGACCTCC	1560
35	TGGTGACTGG AGACAAATGG GACTTTACCC TTCGACTAGA GGATGTGAGC CAGGCCAGG	1620
	CTGGGACCTA CACCTGCCAT ATCCATCTGC AGGAACAGCA GCTCAATGCC ACTGTCACAT	1680
	TGGCAATCAT CACAGTGA CTCCAAATCCT TTGGGTACCT TGGATCCCTG GGAAGCTGC	1740
40	TTTGTGAGGT GACTCCAGTA TCTGGACAAG AACGCTTTGT GTGGAGCTCT CTGGACACCC	1800
	CATCCCAGAG GAGTTTCTCA GGACCTTGGC TGGAGGCACA GGAGGCCCA GCTCCTTTCCC	1860
	AGCCTTGGA ATGCCAGCTG TACCAGGGG AGAGGCTTCT TGGAGCAGCA GTGTACTTCA	1920
	CAGAGCTGTC TAGCCAGGT GCCCAACGCT CTGGGAGAGC CCCAGGTGCC CTCCCAGCAG	1980
45	GCCACCTCCT GCTGTTTCTC ACCCTTGGTG TCCTTTCTCT GCTCCTTTTG GTGACTGGAG	2040
	CCTTTGGCTT TCACCTTTGG AGAAGACAGT GGCAGCAAG ACGATTTTCT GCCTTAGAGC	2100
	AAGGGATTCA CCCTCGCAGG CTCAGAGCAA GATAGAGGAG CTGGAGCAAG AACCGGAGCC	2160
50	GGAGCCGGAG CCGGAACCGG AGCCCGAGCC CGAGCCCGAG CCGGAGCAGC TCTGACCTGG	2220
	AGCTGAGGCA GCCAGCAGAT CTCAGCAGCC CAGTCCAAAT AACGTCCTG TCTAGCAGC	2279

55

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 247 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

15 Leu Gln Pro Gly Ala Glu Val Pro Val Val Trp Ala Gln Glu Gly Ala
 1 5 10 15
 Pro Ala Gln Leu Pro Cys Ser Pro Thr Ile Pro Leu Gln Asp Leu Ser
 20 20 25 30
 Leu Leu Arg Arg Ala Gly Val Thr Trp Gln His Gln Pro Asp Ser Gly
 35 40 45
 Pro Pro Ala Ala Ala Pro Gly His Pro Leu Ala Pro Gly Pro His Pro
 50 55 60
 25 Ala Ala Pro Ser Ser Trp Gly Pro Arg Pro Arg Arg Tyr Thr Val Leu
 65 70 75 80
 Ser Val Gly Pro Gly Gly Leu Arg Ser Gly Arg Leu Pro Leu Gln Pro
 85 90 95
 Arg Val Gln Leu Asp Glu Arg Gly Arg Gln Arg Gly Asp Phe Ser Leu
 100 105 110
 Trp Leu Arg Pro Ala Arg Arg Ala Asp Ala Gly Glu Tyr Arg Ala Ala
 115 120 125
 Val His Leu Arg Asp Arg Ala Leu Ser Cys Arg Leu Arg Leu Arg Leu
 130 135 140
 35 Gly Gln Ala Ser Met Thr Ala Ser Pro Pro Gly Ser Leu Arg Ala Ser
 145 150 155 160
 Asp Trp Val Ile Leu Asn Cys Ser Phe Ser Arg Pro Asp Arg Pro Ala
 165 170 175
 40 Ser Val His Trp Phe Arg Asn Arg Gly Gln Gly Arg Val Pro Val Arg
 180 185 190
 Glu Ser Pro His His His Leu Ala Glu Ser Phe Leu Phe Leu Pro Gln
 195 200 205
 45 Val Ser Pro Met Asp Ser Gly Pro Trp Gly Cys Ile Leu Thr Tyr Arg
 210 215 220
 Asp Gly Phe Asn Val Ser Ile Met Tyr Asn Leu Thr Val Leu Gly Asn
 225 230 235 240
 50 Ser Pro Thr Leu Leu His Ile

245

55

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1629 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

15	TCAGGCTGCC TGATCTGCCC AGCTTTCAG CTTTCTCTG GATTCCGGCC TCTGGTCATC	60
	CCTCCCCACC CTCTCTCCAA GGCCCTCTCC TGGTCTCCCT TCTTCTAGAA CCCCTTCCTC	120
	CACCTCCCTC TCTGCAGAAC TTCTCCTTTA CCCCCACCC CCCACCACTG CCCCTTTTCC	180
20	TTTTCTGACC TCCTTTTGA GGGCTCAGCG CTGCCAGAC CATAGGAGAG ATGTGGGAGG	240
	CTCAGTTCCT GGGCTTGCTG TTTCTGCAGC CGCTTGGGT GGCTCCAGTG AAGCCTCTCC	300
	AGCCAGGGGC TGAGGTCCCG GTGGTGTGGG CCCAGGAGGG GGCTCCTGCC CAGCTCCCCCT	360
	GCAGCCCCAC AATCCCCCTC CAGGATCTCA GCCTTCTGCG AAGAGCAGGG GTCACTTGGC	420
25	AGCATCAGCC AGACAGTGGC CCGCCCGCTG CCGCCCCCGG CCATCCCCTG GCCCCCGGCC	480
	CTCACCCGGC GCGCCCTCC TCCTGGGGGC CCAGGCCCCG CCGCTACACG GTGCTGAGCG	540
	TGGGTCCCGG AGGCCTGCGC AGCGGAGGC TGCCCTGCA GCCCCGCGTC CAGCTGGATG	600
30	AGCGCGGCCG GCAGCGCGGG GACTTCTCGC TATGGCTGCG CCCAGCCCGG CCGCGGACG	660
	CCGGCGAGTA CCGCGCCGCG GTGCACCTCA GGGACCGCGC CCTCTCCTGC CGCCTCCGTC	720
	TGCGCCTGGG CCAGGCCTCG ATGACTGCCA GCCCCCAGG ATCTCTCAGA GCCTCCGACT	780
35	GGGTCATTTT GAACTGCTCC TTCAGCCGCC CTGACCGCCC AGCCTCTGTG CATTGGTTCC	840
	GGAACCGGGG CCAGGCGCGA GTCCCTGTCC GGGAGTCCCC CCATCACCAC TTAGCGGAAA	900
	GCTTCCTCTT CCTGCCCCAA GTCAGCCCCA TGGACTCTGG GCCCTGGGGC TGCATCCTCA	960
40	CCTACAGAGA TGGCTTCAAC GTCTCCATCA TGTATAACCT CACTGTTCTG GGTCTGGAGC	1020
	CCCCAACTCC CTTGACAGTG TACGCTGGAG CAGGTTCCAG GGTGGGGCTG CCCTGCCGCC	1080
	TGCCTGCTGG TGTGGGGACC CGGTCTTTCC TCACTGCCAA GTGGACTCCT CCTGGGGGAG	1140
	GCCCTGACCT CCTGGTGA CTGGAGACAATG GCGACTTTAC CCTTCGACTA GAGGATGTGA	1200
45	GCCAGGCCCC GGCTGGGACC TACACCTGCC ATATCCATCT GCAGGAACAG CAGCTCAATG	1260
	CCACTGTCAC ATTGGCAATC ATCAGAGGTG CCAACGCTC TGGGAGAGCC CCAGGTGCCC	1320
	TCCCAGCAGG CCACCTCCTG CTGTTTCTCA CCCTTGGTGT CCTTCTCTG CTCCTTTTGG	1380
50	TGACTGGAGC CTTTGGCTTT CACCTTTGGA GAAGACAGTG GCGACCAAGA CGATTTTCTG	1440
	CCTTAGAGCA AGGGATTAC CCTCCGACAG CTCAGAGCAA GATAGAGGAG CTGGAGCAAG	1500

55

AACCGGAGCC GGAGCCGGAG CCGGAACCGG AGCCCGAGCC CGAGCCCGAG CCGGAGCAGC 1560
 TCTGACCTGG AGCTGAGGCA GCCAGCAGAT CTCAGCAGCC CAGTCCAAAT AAACGTCCTG 1620
 5 TCTAGCAGC 1629

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 422 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 10
 (ii) MOLECULE TYPE: protein
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
 15
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:
 20 Leu Gln Pro Gly Ala Glu Val Pro Val Val Trp Ala Gln Glu Gly Ala
 1 5 10 15
 Pro Ala Gln Leu Pro Cys Ser Pro Thr Ile Pro Leu Gln Asp Leu Ser
 20 25 30
 Leu Leu Arg Arg Ala Gly Val Thr Trp Gln His Gln Pro Asp Ser Gly
 25 35 40 45
 Pro Pro Ala Ala Ala Pro Gly His Pro Leu Ala Pro Gly Pro His Pro
 50 55 60
 Ala Ala Pro Ser Ser Trp Gly Pro Arg Pro Arg Arg Tyr Thr Val Leu
 65 70 75 80
 30 Ser Val Gly Pro Gly Gly Leu Arg Ser Gly Arg Leu Pro Leu Gln Pro
 85 90 95
 Arg Val Gln Leu Asp Glu Arg Gly Arg Gln Arg Gly Asp Phe Ser Leu
 100 105 110
 35 Trp Leu Arg Pro Ala Arg Arg Ala Asp Ala Gly Glu Tyr Arg Ala Ala
 115 120 125
 Val His Leu Arg Asp Arg Ala Leu Ser Cys Arg Leu Arg Leu Arg Leu
 130 135 140
 40 Gly Gln Ala Ser Met Thr Ala Ser Pro Pro Gly Ser Leu Arg Ala Ser
 145 150 155 160
 Asp Trp Val Ile Leu Asn Cys Ser Phe Ser Arg Pro Asp Arg Pro Ala
 165 170 175
 Ser Val His Trp Phe Arg Asn Arg Gly Gln Gly Arg Val Pro Val Arg
 180 185 190
 45 Glu Ser Pro His His His Leu Ala Glu Ser Phe Leu Phe Leu Pro Gln
 195 200 205
 Val Ser Pro Met Asp Ser Gly Pro Trp Gly Cys Ile Leu Thr Tyr Arg
 210 215 220
 50 Asp Gly Phe Asn Val Ser Ile Met Tyr Asn Leu Thr Val Leu Gly Leu
 225 230 235 240

55

EP 0 900 841 A1

5 Glu Pro Pro Thr Pro Leu Thr Val Tyr Ala Gly Ala Gly Ser Arg Val
 245 250 255
 Gly Leu Pro Cys Arg Leu Pro Ala Gly Val Gly Thr Arg Ser Phe Leu
 260 265 270
 Thr Ala Lys Trp Thr Pro Pro Gly Gly Gly Pro Asp Leu Leu Val Thr
 275 280 285
 10 Gly Asp Asn Gly Asp Phe Thr Leu Arg Leu Glu Asp Val Ser Gln Ala
 290 295 300
 Gln Ala Gly Thr Tyr Thr Cys His Ile His Leu Gln Glu Gln Gln Leu
 305 310 315 320
 15 Asn Ala Thr Val Thr Leu Ala Ile Ile Thr Gly Ala Gln Arg Ser Gly
 325 330 335
 Arg Ala Pro Gly Ala Leu Pro Ala Gly His Leu Leu Leu Phe Leu Thr
 340 345 350
 20 Leu Gly Val Leu Ser Leu Leu Leu Leu Val Thr Gly Ala Phe Gly Phe
 355 360 365
 His Leu Trp Arg Arg Gln Trp Arg Pro Arg Arg Phe Ser Ala Leu Glu
 370 375 380
 25 Gln Gly Ile His Pro Pro Gln Ala Gln Ser Lys Ile Glu Glu Leu Glu
 385 390 395 400
 30 Gln Glu Pro Glu Pro Glu Pro Glu Pro Glu Pro Glu Pro Glu Pro Glu
 405 410 415
 Pro Glu Pro Glu Gln Leu
 420

30 (2) INFORMATION FOR SEQ ID NO: 5:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1468 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: cDNA
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

45 TCAGGCTGCC TGATCTGCCC AGCTTTCAG CTTTCCTCTG GATTCCGGCC TCTGGTCATC 60
 CCTCCCCACC CTCTCTCCAA GGCCCTCTCC TGGTCTCCCT TCTTCTAGAA CCCCTTCTC 120
 CACCTCCCTC TCTGCAGAAC TTCTCCTTTA CCCCCACCC CCCACCACTG CCCCCTTTCC 180
 TTTTCTGACC TCCTTTTGA GGGCTCAGCG CTGCCAGAC CATAGGAGAG ATGTGGGAGG 240
 CTCAGTTCTT GGGCTTGCTG TTTCTGCAGC CGCTTTGGGT GGCTCCAGTG AAGCCTCTCC 300
 50 AGCCAGGGGC TGAGGTCCCG GTGGTGTGGG CCCAGGAGGG GGCTCCTGCC CAGCTCCCCT 360
 GCAGCCCCAC AATCCCCCTC CAGGATCTCA GCCTTCTGCG AAGAGCAGGG GTCACCTGGC 420

55

EP 0 900 841 A1

AGCATCAGCC AGACAGTGGC CCGCCCGCTG CCGCCCCCGG CCATCCCCTG GCCCCCGGCC 480
 CTCACCCGGC GCGGCCCTCC TCCTGGGGGC CCAGGCCCCG CCGCTACACG GTGCTGAGCG 540
 TGGGTCCCGG AGGCCTGCGC AGCGGGAGGC TGCCCCTGCA GCCCCGCGTC CAGCTGGATG 600
 AGCGCGGGCG GCAGCGCGGG GACTTCTCGC TATGGCTGCG CCCAGCCCGG CGCGCGGACG 660
 CCGGCGAGTA CCGCGCCGCG GTGCACCTCA GGGACCGCGC CCTCTCCTGC CGCCTCCGTC 720
 TGCGCCTGGG CCAGGCCTCG ATGACTGCCA GCCCCCAGG ATCTCTCAGA GCCTCCGACT 780
 GGGTCATTTT GAACTGCTCC TTCAGCCGCC CTGACCGCCC AGCCTCTGTG CATTGGTTCC 840
 GGAACCGGGG CCAGGGCCGA GTCCCTGTCC GGGAGTCCCC CCATCACCAC TTAGCGGAAA 900
 GCTTCCTCTT CTGCCCCAA GTCAGCCCCA TGGACTCTGG GCCCTGGGGC TGCATCCTCA 960
 CCTACAGAGA TGGCTTCAAC GTCTCCATCA TGTATAACCT CACTGTTCTG GGTCTGGAGC 1020
 CCCCAACTCC CTTGACAGTG TACGCTGGAG CAGGTTCCAG GGTGGGGCTG CCCTGCCGCC 1080
 TGCCTGCTGG TGTGGGGACC CGGTCTTTCC TCACTGCCAA GTGGACTCCT CCTGGGGGAG 1140
 GCCCTGACCT CCTGGTGACT GGAGACAATG GCGACTTTAC CCTTCGACTA GAGGATGTGA 1200
 GCCAGGCCCA GGCTGGGACC TACACCTGCC ATATCCATCT GCAGGAACAG CAGCTCAATG 1260
 CCACTGTCAC ATTGGCAATC ATCACAGGTC AGCCTCAGGT GGGAAAGGAG TAGCTGCCCT 1320
 CCCAGGGTAG AAAGGACAGG GAGGAAGGGC TGGCAGGGCA AAGACTAGGC AAACCCACCC 1380
 TGTGATGCCA GGCCACTGGG CACAAGTTCC AGAGCCTGCC CATCTCGGCC CCCACTTTTC 1440
 TCACCCCAT AATAAAGAAA CGAAACTG 1468

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 338 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:
 Leu Gln Pro Gly Ala Glu Val Pro Val Val Trp Ala Gln Glu Gly Ala
 1 5 10 15
 Pro Ala Gln Leu Pro Cys Ser Pro Thr Ile Pro Leu Gln Asp Leu Ser
 20 25 30
 Leu Leu Arg Arg Ala Gly Val Thr Trp Gln His Gln Pro Asp Ser Gly
 35 40 45
 Pro Pro Ala Ala Ala Pro Gly His Pro Leu Ala Pro Gly Pro His Pro
 50 55 60

EP 0 900 841 A1

5 Ala Ala Pro Ser Ser Trp Gly Pro Arg Pro Arg Arg Tyr Thr Val Leu
 65 70 75 80
 Ser Val Gly Pro Gly Gly Leu Arg Ser Gly Arg Leu Pro Leu Gln Pro
 85 90 95
 Arg Val Gln Leu Asp Glu Arg Gly Arg Gln Arg Gly Asp Phe Ser Leu
 100 105 110
 10 Trp Leu Arg Pro Ala Arg Arg Ala Asp Ala Gly Glu Tyr Arg Ala Ala
 115 120 125
 Val His Leu Arg Asp Arg Ala Leu Ser Cys Arg Leu Arg Leu Arg Leu
 130 135 140
 15 Gly Gln Ala Ser Met Thr Ala Ser Pro Pro Gly Ser Leu Arg Ala Ser
 145 150 155 160
 Asp Trp Val Ile Leu Asn Cys Ser Phe Ser Arg Pro Asp Arg Pro Ala
 165 170 175
 20 Ser Val His Trp Phe Arg Asn Arg Gly Gln Gly Arg Val Pro Val Arg
 180 185 190
 Glu Ser Pro His His His Leu Ala Glu Ser Phe Leu Phe Leu Pro Gln
 195 200 205
 25 Val Ser Pro Met Asp Ser Gly Pro Trp Gly Cys Ile Leu Thr Tyr Arg
 210 215 220
 Asp Gly Phe Asn Val Ser Ile Met Tyr Asn Leu Thr Val Leu Gly Leu
 225 230 235 240
 30 Glu Pro Pro Thr Pro Leu Thr Val Tyr Ala Gly Ala Gly Ser Arg Val
 245 250 255
 Gly Leu Pro Cys Arg Leu Pro Ala Gly Val Gly Thr Arg Ser Phe Leu
 260 265 270
 35 Thr Ala Lys Trp Thr Pro Pro Gly Gly Gly Pro Asp Leu Leu Val Thr
 275 280 285
 Gly Asp Asn Gly Asp Phe Thr Leu Arg Leu Glu Asp Val Ser Gln Ala
 290 295 300
 40 Gln Ala Gly Thr Tyr Thr Cys His Ile His Leu Gln Glu Gln Gln Leu
 305 310 315 320
 Asn Ala Thr Val Thr Leu Ala Ile Ile Thr Gly Gln Pro Gln Val Gly
 325 330 335
 Lys Glu

- 45 (2) INFORMATION FOR SEQ ID NO: 7:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
TCTCTCAGAG CCTCCGATGG GTCATTTTG 29
5
(2) INFORMATION FOR SEQ ID NO: 8:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
10 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
15
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
TCCTGCAGAT GGATATGGCA GGTGTAGGTC 30
(2) INFORMATION FOR SEQ ID NO: 9:
20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
30 CCTGGGCCAG GCCTCGATGA C 21
(2) INFORMATION FOR SEQ ID NO: 10:
(i) SEQUENCE CHARACTERISTICS:
35 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
40
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
CCCCACTCTG CTTACATTT 20
45
(2) INFORMATION FOR SEQ ID NO: 11:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
50 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:
GGCCCTGGAT CCGGACCTAA TTTTTTTTTT TTTTTT 37

5 (2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
15 GGCCCTGGAT CCGGACCTAA 20

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:
TATAGGATCC GGTGCCCAGA CCATAGGAGA GATG 34

30 (2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:
40 GGCGTTCACG TGGTTGGGCA CCTGTGATGA TT 32

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:
45 (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:
TCACCTACTC GAGAAAAGTG GGGGCCGAGA T 31


55

Claims

1. An isolated nucleotidic sequence selected from the group consisting of :
 - a) the nucleotidic sequences SEQ ID N° 1, SEQ ID N°3 or SEQ ID N°5 ;
 - b) the nucleotidic sequences which hybridize under stringent conditions to any of the sequences defined in a) and which code for a polypeptide which is a variant of the LAG-3 molecule ;
 - c) the nucleotidic sequences which are degenerated as a result of the genetic code to the nucleotidic sequences defined in a) and b), and which code for a polypeptide which is a genetic variant of the LAG-3 molecule.
2. A nucleotidic sequence according to claim 1 selected from the group consisting of the nucleotidic sequences SEQ ID N° 1, SEQ ID N° 3, SEQ ID N° 5 or the fully complementary sequences thereof.
3. A purified polypeptide encoded by a nucleotidic sequence according to claim 1 or 2.
4. A polypeptide according to claim 3, selected from the group consisting of LAG-3V1, LAG-3V2 and LAG-3V3, having a sequence selected from the group consisting of SEQ ID N° 2, SEQ ID N° 4 and SEQ ID N°6 respectively encoded by SEQ ID N° 1, SEQ ID N° 3 or SEQ ID N° 5.
5. An expression vector comprising a nucleotidic sequence according to claim 1 or 2.
6. An host cell transformed with an expression vector according to claim 5.
7. A pharmaceutical composition containing as active ingredient a polypeptide according to claim 3 or 4.
8. Antibodies directed to a specific epitope of one of the polypeptides according to claim 3 or 4.
9. Antibodies according to claim 8, wherein said antibodies are monoclonal antibodies or Fab, Fab', F(ab') or Fv fragments thereof.
10. Use of antibodies according to claim 8 or 9 in a method for purifying, dosing or identifying a polypeptide according to claim 3 or 4.
11. Use of antibodies according to claim 8 or 9 for the manufacture of a therapeutic composition for treating immunerelated pathologies.
12. A therapeutic composition comprising as active ingredient an antibody according to claim 8 or 9.
13. Use of a polypeptide according to claim 3 or 4 for the manufacture of immunomodulators compounds.
14. Use of a polypeptide according to claim 3 or 4 for the manufacture of a therapeutic composition for treating immunerelated pathologies.



FIG. 2






FIG.3





FIG. 3 cont.






FIG.4A




FIG.4B

Lane 1: MW
Lane 2: RT-PCR with primers F459/R460

FIG.5A





FIG.5B



Lane 1: RT-PCR with F176/R460 primers
 Lane 2: Negative control (PCR without RT)
 Lane 3-4: DIG-labelled MW
 Lane 5-6: MW




FIG. 6




FIG. 6
cont.




FIG. 6
cont.




FIG. 7A





FIG. 7B

Lane 1-2: DIG-labelled MW
 Lane 3-4: MW
 Lane 5: RT-PCR with F176/R401 primers
 Lane 6: Negative control (PCR without RT)



Lane 1: RT-PCR with primers 173/V2R
Lane 2: RT-PCR with primers 173/V3R
Lane 3: MW

FIG. 8




FIG. 9



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 97 40 1404

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	WO 91 10682 A (INST NAT SANTE RECH MED ; ROUSSY INST GUSTAVE (FR)) * the whole document *	1-3, 5-14	C12N15/12 C07K14/725 C07K16/28 A61K38/17 A61K39/395 G01N33/53
X	TRIEBEL F ET AL: "LAG-3 A NOVEL LYMPHOCYTE ACTIVATION GENE CLOSELY RELATED TO CD4" JOURNAL OF EXPERIMENTAL MEDICINE, vol. 171, no. 5, 1 May 1990, pages 1393-1405, XP000672284 * the whole document *	1-3, 5, 6	
X	BAIXERAS E ET AL: "CHARACTERIZATION OF THE LYMPHOCYTE ACTIVATION GENE 3-ENCODED PROTEIN. A NEW LIGAND FOR HUMAN LEUKOCYTE ANTIGEN CLASS II ANTIGENS" JOURNAL OF EXPERIMENTAL MEDICINE, vol. 176, no. 2, 1 August 1992, pages 327-337, XP000672285 * the whole document *	3, 7-14	
A	WO 95 30750 A (ROUSSY INST GUSTAVE ; INST NAT SANTE RECH MED (FR); APPLIED RESEARC) * the whole document *	1-14	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C12N C07K A61K G01N
A	WO 88 01304 A (THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK) 25 February 1988 * page 61, line 28 - page 62, line 30 *	1-14	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 November 1997	Examiner Mandl, B
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.92 (Pct/C01)